



Lories and Lorikeets

www.exoticpetvet.co.uk

Lories and lorikeets are colourful parrots, which originate from the islands of South East Asia and Australia. They differ from other parrots by feeding on nectar instead of fruit or seeds and can be found feeding from at least 5,000 species of flowers. To collect the nectar they use their specialised tongues, which have many papillae (extremely fine hairs).

Lories and lorikeets are sexually monomorphic, meaning that they can only be sexed by DNA testing from a blood sample or feather pluck.

Scientific Classification

Lories and lorikeets belong to the Order Psittaciformes, Family Psittacidae, Subfamily Loriinae of which there are 55 different species. One of the most common pets is the colourful Rainbow lorikeet (*Trichoglossus haematodus*).

General care

Ideally these birds will be happiest in a spacious aviary setup. This should be an outdoor enclosure made of strong wire mesh (which should be zinc-free), with an easily cleanable floor and plenty of room for the birds to fly around. In addition to the flying area, a sheltered sleeping area should be provided to protect from the worst of the British weather. The shelter should also provide shade on any sunny days. Perches can be placed at varying heights around the enclosure, with branches of different diameters providing the most natural setup. Appropriate branches can include those from the ash, hazel, birch, willow, eucalyptus, chestnut, sycamore, elder and untreated fruit trees. It is also particularly important to ensure that the enclosure is secure, both to stop the birds getting out (double security doors are the best way to prevent an escape) and to stop predators getting in.

Alternatively, if an aviary is not an option, these birds may be housed indoors in a cage setup. Indoor birds will require more individual attention and stimulation but will often form a close bond with their owners. A cage setup should also be as spacious as possible and placed out of direct sunlight and draughts. It should also be situated away from any item that may give off toxic fumes, such as non-stick cooking utensils, which can release lethal fumes when heated. It is similarly important to ensure that the cage is made out of zinc-free materials as zinc is also very toxic to these birds. A variety of branches can be placed as perches as in an aviary setup, and different toys should be added and changed regularly to entertain birds. If planning to be out of the house for a long time, it is also a good idea to leave a radio program on at a low volume to give the birds some stimulation.

Free flight is an essential requirement and birds should be given the opportunity for exercise daily. It is however, important to ensure that the room they are allowed to fly around is totally secure with all windows, doors and chimneys blocked off, heaters and fans turned off and any potentially poisonous house plants removed or covered. If started young, birds can be easily trained to perch on an arm and returned to the cage.



Barrier Animal Care Clinic LLP

Registered in England Partnership No. OC303416
Members: B Spolander BVSc MRCVS & S Rennie BA VetMB MRCVS
Tel: 020 8293 6580 Fax: 020 8858 5609 Email: wellpetclinic@btconnect.com
www.wellpetclinic.co.uk
VAT Reg: 691 3368 16





Hygiene

Cages should be "spot-cleaned" daily to remove any droppings, feathers, uneaten greens or husks. Once a week, ideally while the birds are exercising, the whole cage and furniture can be disinfected (further details of suitable disinfectants can be obtained from your vet) and branches replaced as necessary.

Diet

In the wild, these birds feed mainly on pollen and nectar from flowers, and various nectar mixes specially designed for lories and lorikeets are available. Please ask for further advice on recommended diets.

Water should be available both for drinking, and in a shallow saucer for bathing.

Fresh food and water should be provided daily and food containers cleaned out.

Lifespan

Lories and lorikeets generally live between 10-20 years.

Signs of health


A healthy bird will be bright and alert with clear eyes and nostrils, shiny feathers and a clean vent. It is important to become familiar with your bird's normal behaviour and droppings, in order that signs of illness can be noticed at an early stage. Beak and nails should also be checked regularly in case trimming is required. It is advisable to take your bird to a vet who routinely deals with birds for a general health check at least once a year.

Signs of illness

Birds will often not show obvious signs of illness until they are very sick, but you should look out for your bird appearing "fluffed up", breathing fast or noisily with its mouth open, any discharges from the eyes or nostrils or any changes in droppings. Changes in beak, nail or feather condition including excessive feather plucking may also indicate a more chronic illness. If you have any concerns, do not hesitate to contact a vet as soon as possible.

This caresheet is only intended as a general guideline, so please ask for further information.

Written and researched by Joanna Hedley BVM&S MRCVS



Barrier Animal Care Clinic LLP
Registered in England Partnership No. OC303416
Members: B Spolander BVSc MRCVS & S Rennie BA VetMB MRCVS
Tel: 020 8293 6580 Fax: 020 8858 5609 Email: wellpetclinic@btconnect.com
www.wellpetclinic.co.uk
VAT Reg: 691 3368 16

